## University of California Agriculture and Natural Resources Making a Difference for California



# UCCE/DWR Weekly Crop Water Use Report

#### WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET<sub>C</sub>) 09/05/25 through 09/11/25

Crops (Leafout Date)	#	#148 Merced			#39 Parlier			#258 Lemon Cove			
	09/05 - 09/11	Accum'd	09/12 - 09/18		09/05 - 09/11	Accum'd	09/12 - 09/18	09/05 - 09/11	Accum'd	09/12 - 09/18	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/1) *	1.48	40.55	1.35		1.46	42.17	1.32	1.40	40.13	1.23	
Pistachio (4/25) * **	1.47	33.27	1.24		1.45	34.77	1.21	1.39	33.42	1.12	
Citrus (2/1)	1.06	32.49	0.98		1.05	34.00	0.96	1.01	32.24	0.88	
Raisin Grapes (4/14) (11 ft. row spacing)	1.02	23.32	0.91		1.00	24.30	0.88	0.95	23.21	0.81	
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis) ***	1.14	25.40	1.03		1.12	26.52	1.00	1.07	25.42	0.93	
Walnuts (4/14)	1.55	32.40	1.24		1.52	33.82	1.21	1.44	32.43	1.15	
Stone Fruit (3/8)	1.55	33.88	1.42		1.53	35.45	1.39	1.47	33.85	1.30	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2025)		0.00				5.40			4.49		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of sarip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

<sup>\*\*</sup> Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

APPLIED WATER		

Crops		#148 Merce	d			#39 Parlier			#258 Lemon Cove				
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/1)	2.3	2.0	1.7	1.6	2.2	1.9	1.7	1.5	2.2	1.9	1.6	1.5	
Pistachio (4/25)	2.3	2.0	1.7	1.5	2.2	1.9	1.7	1.5	2.1	1.9	1.6	1.5	
Citrus (2/1)	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.1	1.6	1.3	1.2	1.1	
Raisin Grapes (4/14) (11 ft. row spacing)***	As	sume all gra	pe	1.1	A	ssume all gra	ipe	1.1	A	1.0			
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis) ***	irrigation type is drip			1.2	irrig	gation type is	drip	1.2	irrig	irrigation type is drip			
Walnuts (4/14)	2.4	2.1	1.8	1.6	2.3	2.0	1.8	1.6	2.2	1.9	1.7	1.5	
Stone Fruit (3/8)	2.4	2.1	1.8	1.6	2.4	2.0	1.8	1.6	2.3	2.0	1.7	1.5	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

### PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops		#148 Merce	ed			#39 Parlier				#258 Lemo	n Cove				
Almonds 115 Trees/A	543	472	401	378	519	449	401	354	519	449	378	354			
Pistachio 106 Trees/A	573	498	424	374	548	473	424	374	523	473	399	374			
Citrus 110 Trees/A	395	346	296	272	395	346	296	272	395	321	296	272			
Raisin Grapes 566 Vines/A	Assume all grape			53	A	ssume all gra	ape	53	A	Assume all grape					
Winegrapes 622 Vines/A	irrig	gation type is	drip	52	irrig	gation type is	drip	52	irrigation type is drip			48			
Walnuts 76 Trees/A	857	750	643	572	822	715	643	572	786	679	607	536			
Stonefruit 172 Trees/A	379	332	284	253	379	316	284	253	363	316	268	237			

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

# UCCE/DWR Weekly Crop Water Use Report





#### WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET<sub>C</sub>) 09/05/25 through 09/11/25

Crops (Leafout Date)	#	#124 Panoche			#2 Five Points			#15 Stratford			
	09/05- 09/11	Accum'd	09/12- 09/18		09/05- 09/11	Accum'd	09/12- 09/18	09/05- 09/11	Accum'd	09/12- 09/18	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (3/1) *	1.52	43.10	1.42		1.62	44.35	1.53	1.60	45.92	1.51	
Pistachio (4/25) * **	1.51	36.05	1.31		1.60	37.02	1.42	1.58	37.98	1.40	
Citrus (2/1)	1.09	35.13	1.00		1.15	36.11	1.11	1.14	37.55	1.09	
Raisin Grapes (4/14) (11 ft. row spacing)	1.04	25.17	0.97		1.09	25.88	1.05	1.08	26.72	1.05	
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.17	27.42	1.09		1.25	28.19	1.18	1.23	29.02	1.16	
Walnuts (4/14)	1.58	35.12	1.31		1.68	35.99	1.42	1.66	37.00	1.40	
Stone Fruit (3/8)	1.59	36.72	1.49		1.68	37.66	1.60	1.66	38.78	1.58	
Past 7 days precipitation (inches)		0.00				0.00			0.00		
Accumulated precipitation (inches) (1/1/2025)		2.32				3.15			2.72		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

<sup>\*\*</sup> Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

DACT WEEKI V	APPLIED WATER	IN INCHES	ADDITION FOR	D FEELCIENCY 1

TAST WEEKET ATTEMED WATER IN INCHES, ADJUSTED FOR EFFICIENCY												
Crops		#124 Panoc	he		#2 Five Points				#15 Stratford			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	2.3	2.0	1.8	1.6	2.5	2.2	1.9	1.7	2.5	2.1	1.9	1.7
Pistachio (4/25)	2.3	2.0	1.8	1.6	2.5	2.1	1.9	1.7	2.4	2.1	1.9	1.7
Citrus (2/1)	1.7	1.5	1.3	1.1	1.8	1.5	1.4	1.2	1.8	1.5	1.3	1.2
Raisin Grapes (4/14) (11 ft. row spacing)	As	sume all gra	pe	1.1	A	ssume all gra	pe	1.1	As	1.1		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.2	irrig	ation type is	drip	1.3	irrig	irrigation type is drip		
Walnuts (4/14)	2.4	2.1	1.9	1.7	2.6	2.2	2.0	1.8	2.6	2.2	2.0	1.7
Stone Fruit (3/8)	2.4	2.1	1.9	1.7	2.6	2.2	2.0	1.8	2.6	2.2	2.0	1.7

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

#### PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops		#124 Panoc	he			#2 Five Poi	nts		#15 Stratford				
Almonds 115 Trees/A	543 472 425		378	590	519	449	401	590	496	449	401		
Pistachio 106 Trees/A	573	498	448	399	623	523	473	424	598	523	473	424	
Citrus 110 Trees/A	420	370	321	272	444	370	346	296	444	370	321	296	
Raisin Grapes 566 Vines/A	As	ssume all gra	ipe	53	A	ssume all gra	ipe	53	Assume all grape			53	
Winegrapes 622 Vines/A	irrig	ation type is	drip	52	irrig	ation type is	drip	57	irrigation type is drip			57	
Walnuts 76 Trees/A	857	750	679	607	929	786	715	643	929	786	715	607	
Stonefruit 172 Trees/A	379	332	300	268	410	347	316	284	410	347	316	268	
For further information concerning all counties receiving this report, contact	the Fresno Co	. Farm Advis	or's office at	(559) 241-75	26.								

<sup>\*</sup> Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.